

Morphology of rice grains recovered from ruins in Thailand

by

Tadayo WATABE* and Tomoya AKIHAMA**

The present paper is a result of morphological analysis of the shape of grains recovered from the bricks collected from old temple-sites and city-walls in central Thailand. Based on our knowledge of the present day methods of brickmaking, the date of these rice grains seems to be identical with that of the bricks. The grains constitute reliable material for identifying the types of rice then cultivated.

I Material and method

The bricks were collected in 1967 from twenty-two different sites shown in Fig. 1. The sites are located in the Mae Nam Chao Phraya basin, an old lowland rice cultivating area. The bricks seem to date from 6th to 18th century, A.D. For the purpose of precise measurement, only the grains in good shape were selected.

Some bricks contained as many as 50 grains each; others only one husk. No grain was found in Sample No. 10. This difference in the number of grains contained (see Table 1) may be attributed to different methods of making the bricks. The bricks containing many grains were probably made of clay mixed with rice hulls—a method still employed in Thailand. Those containing few grains were made perhaps from clay without rice hulls and the grains found in the latter were probably mixed in accidentally. In both cases the bricks bear no vestiges of being fire-baked. They were shaped, and then sun-dried.

II Experimental results and discussion

Results of measurements are shown in Table 1. They show a range of 6.73-7.91 mm in length, 2.89-3.51 mm in breadth, and 2.13-2.62 in length-breadth ratio. Samples Nos. 1 and 2, possibly the oldest grains, are comparatively wide. But the relationship between the morphology and the age of the grains cannot be satisfactorily determined by our data.

* Faculty of Agriculture, Tottori University, Tottori

** National Institute of Agricultural Sciences, Ministry of Agriculture and Forestry, Hiratsuka




Fig. 1 Sampling sites of bricks



Photo. Grain in brick (Collection No. 16)

Rice is usually classified into three types, A, B and C. Type C is *indica*, a variety which is widely cultivated in tropical Asia. Type A is *japonica* commonly grown in the temperate zone. Type B is an intermediate type between A and C. As shown in Fig. 2, almost all varieties of rice cultivated in Thailand belong to type B or C. Only some upland varieties of rice grown by the hill people in northwestern Thailand come on the periphery of type A.*

The results of measurements show that the majority of the grains examined belong to type A. Few exceptions were identified with the other types. We conclude that the varieties of rice cultivated in Thailand in those days differ from the present

* A classification of cultivated rice varieties in Thailand has been proposed by several scientists. Nagamatsu (1942)⁵⁾, Cho (1960)²⁾ and Hamada (1965)³⁾ pointed out that most of the varieties in Thailand belong to *indica*, and Oka (1953)⁶⁾ named them the "continental group" which is composed mainly of the *indica* varieties.

Table Main characters of old grains

No.	Sampling site	Age of sample estimated	Number of grains tested	length mm.	Grain shape breadth mm.	length/ breadth
1	Payuhakhiri	6-11 c A.D.	15	7.91 ± 0.91	3.51 ± 0.11	2.25 ± 0.09
2	Payuhakhiri	6-11 c	39	7.28 ± 0.28	3.41 ± 0.15	2.13 ± 0.10
3	Nakhon Sawan	13-14 c	50	7.45 ± 0.25	3.33 ± 0.18	2.24 ± 0.13
4	Kamphaeng Phet	ca. 14 c	3	6.73 ± 0.55	2.90 ± 0.10	2.32 ± 0.13
5	Kamphaeng Phet	ca. 14 c	30	7.01 ± 0.17	2.90 ± 0.13	2.42 ± 0.08
6	Tha Tako	6-11 c	30	7.36 ± 0.37	3.31 ± 0.20	2.22 ± 0.04
7	Uttaradit	6-11 c	1	6.70	2.90	2.31
8	Sawankhalok	14 c	50	7.29 ± 0.58	3.07 ± 0.18	2.37 ± 0.18
9	Sawankhalok	14 c	50	7.01 ± 0.13	2.99 ± 0.11	2.34 ± 0.09
10	Sawankhalok	14 c	0	---	---	---
11	Sawankhalok	14 c	15	7.31 ± 0.44	3.01 ± 0.22	2.43 ± 0.21
12	Sukhothai	13-15 c	30	7.41 ± 0.67	3.06 ± 0.28	2.42 ± 0.10
13	Sukhothai	13-15 c	30	7.01 ± 0.12	2.95 ± 0.06	2.38 ± 0.07
14	Sukhothai	13-15 c	30	7.15 ± 0.24	3.03 ± 0.16	2.36 ± 0.19
15	Lop Buri	17 c	25	7.10 ± 0.30	3.06 ± 0.26	2.32 ± 0.16
16	Lop Buri	17 c	50	6.94 ± 0.16	3.23 ± 0.16	2.15 ± 0.10
17	Ayutthaya	14-18 c	30	7.50 ± 0.38	3.13 ± 0.17	2.40 ± 0.09
18	Ayutthaya	14-18 c	30	7.06 ± 0.22	3.13 ± 0.25	2.26 ± 0.18
19	Nan	?	1	6.60	2.80	2.36
20	Phrae	?	1	7.40	2.80	2.64
21	Phrae	?	5	7.20 ± 0.25	3.00 ± 0.07	2.40 ± 0.07
22	Sing Buri	?	50	7.56 ± 0.23	2.89 ± 0.14	2.62 ± 0.12

lowland rice varieties. They show much similarity to those varieties now cultivated in the upland area of Thailand. It is to be noted, however, that the present physiological conditions of the sampling sites are unsuitable for the upland varieties.

The transition of grain type from *japonica* to *indica* in a period of only two hundred years can hardly be accepted as the result of botanical evolution. It may be attributed to a change caused by external reasons. A sudden increase of rice production for export after the Bowring Treaty of 1855 came into effect may have led to the breeding or introduction of new varieties.

In the present paper only the morphological aspects are taken up. Chemical analysis and determination of date by the C₁₄ dating method are now being done to confirm our result.

Acknowledgements

The authors would like to express their hearty thanks to those of NRC of Thailand—General Netr Khemayodhin, Dr. Pradisth Cheosakul, Dr. Prasart Na Nagara and Mrs. Boonthom Thamcharee—for their assistance and support. The authors




Fig. 2 Distribution of grain shape, data by Akihama (1965)¹⁾ and Watabe (1967)⁷⁾ are plotted on the Matsuo's classification map (1952)⁴⁾

are indebted to Professor Shinobu Iwamura, former Director of the Center, for valuable advice and encouragement, and to Professor Yoneo Ishii and Dr. Yoshikazu Takaya of the Center for their cooperation and helpful discussions. We also appreciate Mr. Victor Kennedy, Medical University of Bangkok, who kindly collected some of the specimens for the present study.

References

- 1) Akihama, T. "Classification of Cultivated Rice Varieties," *Nature and Life in Southeast Asia*, 4: 88-95. Osaka, 1965.
- 2) Cho, J. *Cultivated Rice Plant in the Basin of Mekong River*. Tokyo: Jap. Soc. Ethnology, 1960.
- 3) Hamada, H. "Rice in the Mekong Valleys," *Indo-Chinese Studies; Synthetic Research of the Culture of Rice-cultivating Races in Southeast Asian Countries*, 1: 507-586. Tokyo, 1965.
- 4) Matsuo, T. "Genecological Studies on the Cultivated Rice," (Jap.), *Natl. Inst. Agr. Sci. Ser.*, D 3: 1-111. Tokyo, 1952.
- 5) Nagamatsu, T. "Studies on the Geographical Differentiation of Cultivated Rice. III," (Jap.), *Proc. Crop Sci. Soc. Japan*, 14: 132-145. Tokyo, 1942.
- 6) Oka, H. "Phylogenetic Differentiation of the Cultivated Rice Plant, I," *Jap. J. Breed.*, 3: 33-43. Tokyo, 1953.
- 7) Watabe, T. *Glutinous Rice in Northern Thailand*. Kyoto: Kyoto University, 1967.